



Rec'd PCT/PTO

24 JAN 2005 #2



INVESTOR IN PEOPLE

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

EP/03/8197

The Patent Office
Concept House
Cardiff Road
Newport

South Wales
NP10 8QQ


REC'D 05 NOV 2003
WIPO PCT

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

stration under the Companies Act does not constitute a new legal entity but merely the company to certain additional company law rules.




D. Mahoney

Signed

Dated 6 August 2003

Patent of 1977
(Rule 1a)



1/77
29 JUL 02 E736571-2 002093
P01/7700 0.00-0217406.8
The Patent Office

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

Cardiff Road
Newport
South Wales
NP10 8QQ

1. Your reference

PPD70059/GB

0217406.8

26 JUL 2002

3. Full name, address and postcode of the or of each applicant (*underline all surnames*)

SYNGENTA PARTICIPATIONS AG
Intellectual Property Department
Schwarzwaldallee 215
4058 Basel
SWITZERLAND

Patents ADP number (*if you know it*)

If the applicant is a corporate body, give the country/state of its incorporation

8029555001

4. Title of the invention

DISEASE RESISTANT BRASSICAS

5. Name of your agent (*if you have one*)

"Address for service" in the United Kingdom to which all correspondence should be sent (*including the postcode*)

JOZSEF CHRISTOPHER GAAL
Intellectual Property Department
Syngenta Limited
Jealott's Hill International Research Centre
PO Box 3538
Bracknell, Berkshire, RG42 6YA
UNITED KINGDOM

Patents ADP number (*if you know it*)

7991854002

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (*if you know it*) the or each application number

Country

Priority application number
(*if you know it*)

Date of filing
(*day / month / year*)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing
(*day / month / year*)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (*Answer 'Yes' if:*

- a) any applicant named in part 3 is not an inventor, or
 - b) there is an inventor who is not named as an applicant, or
 - c) any named applicant is a corporate body.
- See note (d))

YES (b)

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description 20

Claim(s) 04

Abstract 01

Drawing(s) 00

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (*Patents Form 7/77*)

Request for preliminary examination and search (*Patents Form 9/77*)

Request for substantive examination (*Patents Form 10/77*)

Any other documents
(please specify)

11.

I/We request the grant of a patent on the basis of this application.
SYNGENTA PARTICIPATIONS AG

Signature *Joanna Chandler* Date *25/7/02*
Authorised Signatory

12. Name and daytime telephone number of person to contact in the United Kingdom

Joanna Carmen CHANDLER = 01344 414365

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- a) If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505.
- b) Write your answers in capital letters using black ink or you may type them.
- c) If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- d) If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- e) Once you have filled in the form you must remember to sign and date it.
- f) For details of the fee and ways to pay please contact the Patent Office.

DISEASE RESISTANT BRASSICAS

The present invention relates to plants resistant to diseases, in particular to *Brassica oleracea* plants resistant to the disease clubroot.

5 Clubroot is a widespread disease that causes serious problems in many Brassica growing areas (for a review see e.g. Dixon (1999) Grower April 29, pp.28-29). The disease is caused by *Plasmodiophora brassicae*, a one-cellular organism. Symptoms of the disease include root malformations with hard swellings (clubs), that eventually rot. The disease also causes stunting through reduced growth, and wilting of leaves is observed under water stress.
10 Chemical control of the disease is not effective. Therefore, good genetic resistance of the crop is important for its protection against the disease.

The genus of the Brassicas comprises several species of commercial interest, such as *B. rapa* (Chinese cabbage, pak choi, turnip), *B. napus* (oil seed, swede), *B. juncea* (mustard), *B. nigra* (black mustard) and *B. oleracea* (cauliflower, broccoli, cabbage, Brussels sprouts,
15 savoy cabbage, borecole, kohlrabi, borecole and others). While sub-species within a species of the Brassica genus are usually sexually compatible, this is not necessarily the case between different species of the Brassica genus. For example, *B. rapa* and *B. oleracea* do not have the same number of chromosomes (10 chromosomes versus 9 chromosomes) and are therefore not sexually compatible. This renders the transfer of a trait from one Brassica
20 species to another particularly difficult.

Several sources of resistance to clubroot have been described within the Brassica genus (see e.g. Bradshaw et al. (1997) Ann. Appl. Biol. 130:337-348; Gowers (1982) Euphytica 31:971-976). Some resistances are monogenic, some polygenic, some are dominant, some recessive. Monogenic dominant resistances have been described in *B. rapa*
25 and *B. napus*, such as for example a monogenic dominant resistance in the *B. rapa* Chinese cabbage (Yoshikawa (1983) Japan Agricultural Research Quarterly, Vol.17, no.1, p. 6-11). Chinese cabbage F1-hybrids with this resistance have been shown to have good protection against clubroot, although a small number of strains ('races') of clubroot have been able to break through this resistance, especially under warm conditions (temperatures above ca. 25
30 °C). Such races seem more prevalent in Asia than in Europe.

By contrast, only polygenic, recessive sources of resistance have been described in the Brassica species *B. oleracea* (see e.g. Voorrips (1995) Euphytica 83:139-146). Such

sources have proven not only to be insufficiently resistant to clubroot, but they are also very difficult to transfer between to commercial *B. oleracea* lines. This renders the breeding of the resistance a difficult and time consuming task.

Therefore, there is an unfulfilled need for *B. oleracea* plants with an improved resistance to clubroot, wherein the resistance is also easy to breed and transfer to commercial *B. oleracea* lines.

Accordingly, the present invention addresses the problem of unsatisfactory resistance to the disease clubroot in *B. oleracea*. To achieve improved resistance to the disease in *B. oleracea*, the present invention discloses the transfer of a monogenic dominant resistance to clubroot from chinese cabbage (*B. rapa*) to the *B. oleracea* broccoli, and then further to other *B. oleraceas*, such as white cabbage, cauliflower and Brussels sprouts. In order to overcome the sexual incompatibility between *B. rapa* and *B. oleracea*, the resistance to clubroot was transferred by means of interspecific hybridization using an embryo-rescue technique, followed by repeated backcrosses and disease tests in all backcross generations. High level of resistance to clubroot is obtained in the resulting *B. oleracea* plants. The resistance is stable, and can be transmitted to further generations and transferred to susceptible or less resistant *B. oleracea* plants.

Thus, the present invention discloses *B. oleracea* plants resistant to clubroot, wherein the resistance to clubroot is monogenic and dominant, including seeds and materials of said plants and the progeny thereof. The present invention also discloses methods to produce *B. oleracea* plants resistant to clubroot, methods to transfer the clubroot resistance to susceptible or less resistant *B. oleracea* plants. The present invention also further discloses molecular markers linked to the resistance to clubroot.

The present invention is particularly advantageous over existing resistances to clubroot in *B. oleracea* in that the resistance of the instant invention is easily transferred between *B. oleracea* plants and commercial lines. Higher yields are obtained because of the absence of disease on resistant plants. Moreover much less crop protection chemicals or no crop protection chemicals at all are required against clubroot when *B. oleracea* plants of the present invention are grown.

The present invention therefore discloses:

A *B. oleracea* plant resistant to clubroot, wherein the resistance to clubroot is monogenic and dominant. Preferably, the *B. oleracea* plant is rated at level 2 or less in a test for the disease having a 1-9 scale. Preferably, the *B. oleracea* plant is rated at level 1 in a test for the disease having a 1-9 scale. Preferably, the *B. oleracea* plant is rated at level 1 or less in a test for the disease having a 0-5 scale. Preferably, the *B. oleracea* plant is rated at level 1 or less in a test for the disease having a 0-5 scale. In another preferred embodiment, the *B. oleracea* plant is broccoli, white cabbage, cauliflower, Brussels sprouts, Borecole, Savoy, or red cabbage. In another preferred embodiment, the *B. oleracea* plant is homozygous or heterozygous for the resistance to clubroot. In another preferred embodiment, the resistance to clubroot is linked to a molecular marker obtainable by PCR amplification using primer O20 (SEQ ID NO:1) or primer Y13 (SEQ ID NO:2). Preferably, the resistance is within 10 cM of said molecular marker, preferably within less than 6 cM, preferably within less than 5 cM, more preferably within less than 3 cM, even more preferably within less than 1.5 cM. In another preferred embodiment, the resistance is obtainable from a clubroot resistant *B. rapa* plant, preferably from Chinese cabbage F1 hybrid Parkin. In another preferred embodiment, the plant is an inbred or a hybrid. In yet another preferred embodiment, the plant is cytoplasmic male sterile. In another preferred embodiment, the present invention discloses a *B. oleracea* plant comprising a locus conferring a monogenic, dominant clubroot resistance to said plant. Preferably, the locus is within 10 centi-Morgan (cM) of a molecular marker obtainable by PCR amplification using primer O20 (SEQ ID NO:1) or primer Y13 (SEQ ID NO:2), preferably within 6 cM, preferably within 5 cM, more preferably within 3 cM, even more preferably within 1.5 cM of said molecular marker. In another preferred embodiment, the resistance to clubroot according to the present invention is obtainable from line CFL667, deposited with NCIMB under accession number NCIMB 41134.

The present invention further discloses:

Seed of a plant disclosed above, including the progeny thereof.

Fruit of a plant disclosed above. Part of a plant disclosed above.

The present invention further discloses:

Use of a monogenic and dominant resistance to clubroot to confer a *B. oleracea* plant resistance to said disease, wherein said resistance is obtainable from a *B. rapa* plant, preferably from Chinese cabbage F1 hybrid Parkin.

5 The present invention further discloses:

A method for producing a *B. oleracea* plant comprising a monogenic and dominant resistance to clubroot comprising the steps of: a) obtaining a *B. rapa* plant resistant to clubroot; b) crossing said *B. rapa* plant with a *B. oleracea* plant, c) rescuing embryos resulting from the cross of step b); d) regenerating a plant from a embryo of step c); e) 10 selecting a plant of step d) that is resistant to clubroot; f) back-crossing a plant resulting from step e) with a *B. oleracea* plant. In a preferred embodiment, the method further comprises introgressing the resistance into an elite *B. oleracea* inbred. In another preferred embodiment, the method further comprises crossing said inbred to another *B. oleracea* inbred to produce a hybrid.

15

The present invention further discloses:

A *B. oleracea* plant obtainable by a method disclosed above.

The present invention further discloses:

20 A method for transferring a monogenic and dominant resistance to clubroot to a *B. oleracea* plant susceptible or less resistant to the disease comprising the steps of: a) obtaining a *B. oleracea* plant comprising a monogenic and dominant resistance to clubroot; b) crossing said *B. oleracea* plant of step a) with a *B. oleracea* plant susceptible or less resistant to clubroot; c) selecting a plant from the cross of step b) that is resistant to clubroot. In a preferred 25 embodiment, the method further comprises backcrossing said resistance into said *B. oleracea* plant susceptible or less resistant to clubroot.

The present invention further discloses:

30 A DNA fragment amplified from a Brassica genome, wherein said DNA fragment is approximately 400 bp long and comprises SEQ ID NO:1.

A DNA fragment amplified from a Brassica genome, wherein said DNA fragment is approximately 640 bp long and comprises SEQ ID NO:2.

In a preferred embodiment, a DNA fragment disclosed above is indicative of the presence of a dominant and monogenic resistance to clubroot in a Brassica plant.

Use of a DNA fragment disclosed above to identify a Brassica plant that is resistant to clubroot.

- 5 Use of primer O20 (SEQ ID NO:1) to detect a DNA fragment of approximately 400 bp in a Brassica genome.

Use of primer Y13 (SEQ ID NO:2) to detect a DNA fragment of approximately 640 bp in a Brassica genome.

Use of a primer O20 or Y13 to identify a Brassica plant resistant to clubroot.

- 10 In a preferred embodiment, the Brassica plant is *B. oleracea*.

A kit for detecting a monogenic and dominant resistance to clubroot in a *B. oleracea* plant comprising an oligonucleotide of set forth in SEQ ID NO:1 or SEQ ID NO:2.

The present invention further discloses:

- 15 A method for transferring a monogenic and dominant resistance to clubroot to a *B. oleracea* plant susceptible or less resistant to the disease comprising the steps of: a) obtaining a *B. oleracea* plant comprising a monogenic and dominant resistance to clubroot; b) crossing said *B. oleracea* plant of step a) with a *B. oleracea* plant susceptible or less resistant to clubroot; c) selecting a plant comprising a DNA fragment obtainable by PCR amplification using
20 primer O20 (SEQ ID NO:1) or primer Y13 (SEQ ID NO:2), wherein said plant of step c) is resistant to clubroot. In a preferred embodiment, the method further comprises backcrossing said resistance into said *B. oleracea* plant susceptible or less resistant to clubroot.

The present invention further discloses:

- 25 A method of identifying a *B. oleracea* plant resistant to clubroot comprising the steps of: a) obtaining a sample from a *B. oleracea* plant; b) detecting in said sample a DNA fragment obtainable by PCR amplification using primer O20 (SEQ ID NO:1) or primer Y13 (SEQ ID NO:2), wherein said *B. oleracea* plant of step b) is resistant to clubroot.

- 30 The present invention further discloses:

A method of selecting a *B. oleracea* plant resistant to clubroot from a population of *B. oleracea* plants comprising the steps of: a) providing a population of *B. oleracea* plants; b)

obtaining a sample of a plant of said population; c) detecting in said sample a DNA fragment obtainable by PCR amplification using primer O20 (SEQ ID NO:1) or primer Y13 (SEQ ID NO:2), wherein said *B. oleracea* plant of step b) is resistant to clubroot.

5

Definitions

Trait: characteristic or phenotype, for example a resistance to a disease. A trait may be inherited in a dominant or recessive manner, or may be monogenic or polygenic. A trait is for example a resistance to a disease, such as clubroot.

Resistance: characteristic or phenotype of a plant to exhibit no symptoms or insignificant symptoms of a disease. Resistance is: thus preferably the ability of a plant to reduce the development of a pathogen. (see for example Robinson, R.A., 1969: Review Applied Mycology 48, 593-606).

Monogenic: determined by a single locus.

Polygenic: determined by more than one locus.

15 Dominant: determines the phenotype when present at the heterozygous or homozygous state.

Recessive: only displayed when present at the homozygous state.

Locus: region on a chromosome, which comprises one or more genetic factors, for example one or several genes, contributing to a trait, such as a resistance to a disease.

20 Genetic linkage: tendency of chromosomal regions to be inherited together as a result of their location on the same chromosome. Measured by percent recombination between loci (centi-Morgan, cM).

Isogenic: plants which are genetically identical, except that they may differ by the presence or absence of a heterologous DNA sequence.

25 Marker assisted selection: refers to the process of selecting a desired trait or desired traits in a plant or plants by detecting one or more nucleic acids from the plant, where the nucleic acid is associated with the desired trait.

The present invention discloses *B. oleracea* plants resistant to the disease clubroot. In particular, the present invention discloses *B. oleracea* plants, which comprise a
30 monogenic, dominant resistance to the disease. This resistance provides improved resistance

to the disease, when compared to existing resistances in *B. oleracea* and can be easily transferred between *B. oleracea* plants.

*B. oleracea*s plants of the present invention are for example:

Brassica oleracea L. (cole-crops)

- 5 • var. *acephala* DC. (kales)
- var. *albiflora* Sun [= *B. alboglabra*] (Chinese kale)
- var. *alboglabra* [= *B. alboglabra*] (Chinese kale)
- var. *botrytis* L. (cauliflower, heading broccoli)
- var. *capitata* L. (cabbage)
- 10 • var. *chinensis* Prain (burma sarson)
- var. *fimbriata* Mill. (kitchen kale)
- var. *fruticosa* Metz. (thousand-head kale)
- var. *gemmifera* DC. (brussels sprouts)
- var. *gongylodes* L. (kohlrabi)
- 15 • var. *italica* Plenck. (broccoli, calabrese)
- var. *sabauda* L. (savoy cabbage)
- var. *sabellica* (collards)
- var. *tranchuda* L.H. Bailey (tranchuda cabbage)

Preferred *B. oleracea* plants of the present invention are white cabbage, cauliflower, Brussels
20 sprouts and broccoli.

According to the present invention a monogenic dominant resistance to clubroot is transferred from chinese cabbage (*B. rapa*) to broccoli (*B. oleracea*). Interspecific hybridization was carried out between broccoli (*B. oleracea* var. *italica*) as female parent and *B. rapa* as male parent. Broccoli inbred lines were chosen as female to prevent cell
25 organelles of different origins in the end products of the cross. The male parent in the interspecific cross was a clubroot resistant Chinese cabbage F1 hybrid from Japan, commercially available under the name "Parkin" (Takii Seeds, Japan). After each cross or back-cross, chromosomes of the resulting plants were counted to assess the level of ploidy of the plants and the progress towards obtaining a *B. oleracea* chromosome set. Chromosome
30 counting was carried out according to the method of Chiang et al. (1979) Euphytica 28: 41-45. Flow cytometer work was done with a Partec CA II (Partech, UK).

F1 hybrids resulting from the cross were obtained by embryo rescue. Embryo rescue had to be used since *B. rapa* and *B. oleracea* do not have the same number of chromosomes and are therefore not sexually compatible. Embryo rescue allows to overcome the problems of degradation of the endosperm. The embryo rescue technique described by Harbert et al. (Euphytica 18 (1969) p. 425-429) was used. Sterilized ovules of 10-12 days old after pollination are cut in half and introduced to a culture medium kept in constant motion. The culture medium is basically White's medium with 8% sucrose and 400 ppm caseine hydrolysate. The embryo's are washed out of the ovules and grow in the liquid medium. Clubroot resistant F1 plants were crossed with broccoli (i.e. backcrossed). Embryo rescue as described above was also required to obtain plants resulting from the first backcross (i.e. BC1 plants).

BC1 plants resistant to clubroot were backcrossed again with broccoli and three clubroot resistant BC2 plants were obtained by normal seed set (designated B27, B23 and B28). Plant B27 was further backcrossed with broccoli and plants resulting from the fourth back-cross (BC4) of plant B27 were used as source of the resistance for other *B. oleracea* crops.

The detailed experimental protocol leading to the transfer of the clubroot resistance to *B. oleracea* is described in Example 1. Disease tests used to test for the presence of the resistance are described in Example 2. Results of field trials demonstrating resistance to clubroot are shown in Example 3.

A representative *B. oleracea* line comprising the monogenic dominant resistance to clubroot according to the present invention, line CFL667, was deposited with NCIMB, Aberdeen AB2 1RY, Scotland, UK under accession number NCIMB 41134 on June 28, 2002. Accordingly, the resistance to clubroot according to the present invention is obtainable from line CFL667, deposited with NCIMB under accession number NCIMB 41134.

In a preferred embodiment of the present invention, a *B. oleracea* plant resistant to clubroot is defined as a plant assessed at rating 0 or 1 in the 0-5 scale of example 2 herein, or as a plant assessed at rating 1 or 2 in the 1-9 scale of example 2 herein.

The resistance to clubroot was transferred to other *B. oleraceas*, in particular white cabbage, cauliflower and Brussels sprouts, using standard breeding techniques in the Brassica art. The trait was also further introgressed into *B. oleracea* elite lines. The introgression of the resistance into the elite line is for example achieved by recurrent

selection breeding, for example by backcrossing. In this case, the elite line (recurrent parent) is first crossed to a donor inbred (the non-recurrent parent) that carries the resistance. The progeny of this cross is then mated back to the recurrent parent followed by selection in the resultant progeny for clubroot resistance. After three, preferably four, more preferably five or more generations of backcrosses with the recurrent parent with selection for clubroot resistance, the progeny is heterozygous for the locus harboring the resistance, but is like the recurrent parent for most or almost all other genes (see, for example, Poehlman & Sleper (1995) *Breeding Field Crops*, 4th Ed., 172-175; Fehr (1987) *Principles of Cultivar Development*, Vol. 1: Theory and Technique, 360-376, incorporated herein by reference).

Selection for clubroot resistance is carried out after each cross. In a preferred embodiment, the present invention therefore encompasses a method of transferring a dominant and monogenic resistance to clubroot into *B. oleracea* plant susceptible or less resistant to the disease comprising the following steps: crossing a first *B. oleracea* plant with a second *B. oleracea* plant, wherein the first plant comprises a dominant and monogenic resistance to clubroot, selecting a plant resulting from the cross that is resistant to clubroot. The method further comprises backcrossing a plant selected above with the second *B. oleracea* plant until the resistance has been transferred to the second *B. oleracea* plant.

The clubroot resistance is preferably transferred to commercial lines of *B. oleracea*. Such lines are for example inbred lines. Alternatively, commercial lines are hybrids, which are obtained by crossing two inbred lines. In this case, the resistance to clubroot may be present in one of the parent inbreds or in both. Therefore, in a preferred embodiment, the present invention discloses an inbred or a hybrid line of *B. oleracea* comprising a monogenic dominant resistance to clubroot, including seeds and materials of said inbred or hybrid and the progeny thereof. Preferably, the resistance is obtainable from a *B. rapa*, preferably from Chinese cabbage F1 hybrid "Parkin". In another preferred embodiment, the resistance to clubroot according to the present invention is present in line CFL667, deposited with NCIMB under accession number NCIMB 41134.

In another preferred embodiment, a clubroot resistance *B. oleracea* plant of the present invention is male sterile. Male sterility is of value in *B. oleracea* hybrid seed breeding because normal flowers are self-pollinating. Male sterile lines do not produce viable pollen and cannot self-pollinate. By eliminating the pollen of one parental variety in a cross, a plant breeder is assured of obtaining hybrid seed of uniform quality. A particularly

advantageous male sterility system is cytoplasmic male sterility (CMS). An example of such CMS in Brassica is the Ogura CMS originally found in radish (see for example Ogura (1968) Mem. Fac. Agric. Kagoshima Univ. 6: 39-78; Makaroff (1989) Journal of Biol. Chem. 264: 11706-11713; US 5,254,802). Therefore, the present invention discloses a male sterile, in
5 particular CMS, *B. oleracea* plant comprising a monogenic dominant resistance to clubroot, including seeds and materials of said plants and the progeny thereof. Preferably, the resistance is obtainable from *B. rapa*, preferably from Chinese cabbage F1 hybrid "Parkin". Preferably, the CMS is derived from an Ogura genome.

In yet another preferred embodiment, the present invention discloses molecular
10 markers, which are linked to the monogenic dominant resistance to clubroot. Such molecular markers allow to differentiate at the molecular level between resistant and susceptible plants, and are thus indicative of the monogenic and dominant resistance in Brassica plants, preferably *B. oleracea* plants. These molecular markers are based on genetic polymorphisms between resistant and susceptible plants and are located at the locus of the resistance or
15 closely nearby. Molecular markers are for example used to breed and obtain resistant plants, to follow the presence of the resistance during breeding, or to control the presence of the resistance in a commercial seed lot.

Thus, in a preferred embodiment, the present invention provides methods for mapping the clubroot resistance in Brassica plants, preferably *B. oleacea* plants, methods for
20 determining whether the resistance is present in a Brassica plant, preferably a *B. oleacea* plant, and methods for transferring the resistance to a susceptible or less resistant Brassica plant, preferably a *B. oleacea* plant. For example, the methods are used in breeding new resistant plants or lines or to control the quality of a seed lot. The molecular markers disclosed herein allow for quicker release of resistant lines to the market and for a better
25 quality control of commercial seed lots. A disease assay can thus be avoided, and replaced by the use of a molecular marker.

The molecular markers and methods of the present invention are particularly beneficial in the various steps leading to a commercial variety, such as in breeding programs and in the seed production process. For example, the present invention is used to introgress
30 the resistance into a *B. oleracea* plant, for example an elite inbred line. Selection for the resistance to clubroot is carried out by testing seeds or plants resulting from each cross with molecular markers. In a preferred embodiment, the present invention therefore encompasses

methods of transferring a dominant and monogenic resistance to clubroot into a *B. oleracea* plant susceptible or less tolerant to the disease comprising the following steps: crossing a first plant resistant to the disease with a second plant susceptible or less resistant, harvesting the seeds resulting from the cross, obtaining a sample of the seed or of a plant grown therefrom, detecting in said sample a molecular marker of the present invention, the presence of said molecular marker being indicative of the resistance to the disease in said seed or plant, selecting a plant resistant to the disease. Preferably, the resistance is then further backcrossed into the susceptible or less resistant plant.

Molecular markers of the present invention are also conveniently used in the production of stable homogenous inbred lines or cultivars (also sometimes called varieties), whereby a particular line is self-pollinated until satisfactory purity and homogeneity of the line is reached. The present invention is similarly used for the commercial production of seeds of a particular line or cultivar. Here again, after each cross a method of the present invention is applied to the seeds resulting from the cross and only resistant plants are selected. In a preferred embodiment, the present invention therefore encompasses methods of producing clubroot resistant seeds or plants comprising the following steps: crossing a resistant plant with itself, harvesting the seeds resulting from the cross, obtaining a sample of the seed or of a plant grown therefrom, detecting in said sample a molecular marker of the present invention, the presence of said molecular marker being indicative of the resistance to the disease in said seed or plant.

Similarly, the present invention is used in hybrid seed production. In this case, the present invention is used to assure that all hybrid seeds that germinate and grow in the field are resistant to the disease. Preferably, molecular markers of the present invention are also used in quality assurance to ensure that the disease resistance is present in the hybrid seeds. In a preferred embodiment, the present invention therefore encompasses methods producing seeds comprising the following steps: crossing a first plant with a second plant, wherein one of the plants is resistant to clubroot, harvesting the seeds resulting from the cross, obtaining a sample of the seed or of a plant grown therefrom, detecting in said sample a molecular marker of the present invention, the presence of said molecular marker being indicative of the resistance to the disease in said seed or plant. The present invention thus provides a significant advancement to commercial breeding and seed production processes.

In a preferred embodiment, a molecular marker of the present invention is within 10 centi-Morgan (cM) from the locus of the clubroot resistance, preferably within 6 cM, preferably within 5 cM, more preferably within 3 cM, even more preferably within 1.5 cM from the locus of the clubroot resistance, and thus tightly co-segregates with the resistance.

5 Preferred polymorphisms in the context of the present invention include for example single sequence repeats (SSR, see e.g. Hearne et al. (1992) Trends Genet 8:288-294), single nucleotide polymorphisms (SNP, Botstein B et al. (1980) Am J Hum Genet 32:314-331), and any other types of DNA rearrangements, such as deletions, amplifications, crossovers.

Methods for detection of polymorphisms well known in the art can be applied in the
10 context of the present invention to produce a molecular marker linked to the resistance. Preferred methods include methods based on the PCR amplification technique, for example the SSR technology and the RAPDs technology (Williams et al. (1990) Nucl Acids Res 18:6531-6535). In general, preferred oligonucleotides for PCR are about 8 to 50 nt long, more preferably 10 to 30 nt long. PCR amplified DNA fragments comprising a site
15 comprising a polymorphism of the present invention are preferably about 100-3,000 nt long, more preferably about 200 to 2,000 nt long, even more preferably about 300 to 1,000 nt long. Other methods for screening or detecting polymorphisms, which are applicable to the instant invention include direct sequencing of nucleic acids, single strand polymorphism assay, ligase chain reaction, enzymatic cleavage, and southern hybridization.

20 Alternative methods include several methods for detecting Cleaved Amplified Polymorphic Sequences (CAPS; Konieczny et al., The Plant Journal 4(2):403-410, 1993) and for detecting Single Nucleotide Polymorphisms (SNPs) with a method termed "CAMPS" for Cleaved Amplified Modified Polymorphic Sequences, also known as dCAPS (Neff et al, 1998. Plant J. 14: 387-392; Michaels and Amasino, 1998. Plant J 14: 381-385). In the CAPS
25 method, a nucleic acid containing a polymorphic restriction site is amplified using primers flanking the restriction site. The resulting PCR product is digested with the restriction endonuclease corresponding to the polymorphic restriction site, and the digested products are analyzed by gel electrophoresis.

Southern hybridization is also an effective method of identifying differences in
30 sequences, in particular using the RFLP (restriction fragment length polymorphisms) technology (Botstein B et al. (1980) Am J Hum Genet 32:314-331).

In a preferred embodiment, the present invention discloses molecular markers closely linked to the clubroot resistance obtained using the RAPDs technology (see example 4 herein).

5 The following examples are intended to provide illustrations of the application of the present invention. The following examples are not intended to completely define or otherwise limit the scope of the invention.

Examples

Example 1: Transfer of the resistance to *B. oleracea*

10 F1

A few hundred crosses between the clubroot resistant Chinese cabbage and broccoli were performed. From these crosses, embryos were rescued. Four of these embryos developed into plants, two of which proved to be resistant to the disease. These two resistant plants had 19 chromosomes. The phenotype was a hybrid between Chinese cabbage and broccoli.

15 BC1

Embryo rescue was conducted again to obtain BC1 plants. Broccoli inbred lines were used as backcross parent. Broccoli lines were also used a backcross parent for the following generations. Only one of the two F1 plants mentioned above responded and 5 embryos developed into plants. Four of them were resistant to clubroot.

20 The phenotype of the BC1 plants was closer to that of the broccoli backcross parent. As the distribution of chromosomes of the *B. rapa* genome in the BC1 plants is at random, the plants contain 18 chromosomes of *B. oleracea* and 0-10 chromosomes of *B. rapa*. Accordingly, plants with 18-28 chromosomes were observed.

BC2

25 BC2 plants were obtained by seed setting on the BC1 plants. Nine plants originating from seedset on a plant designated as plant B from the BC1 above proved to be resistant to clubroot. The number of chromosomes was determined and the deviation from the normal DNA content was determined with a flowcytometer. Three plants, B23, B27, and B28 were further use in the program.

BC3 and BC4

BC3 and BC4 were also obtained by seed set. Plant B27, which was already diploid in the BC2, had incorporated the dominant clubroot resistance gene. The backcross generations BC3 and BC4 show the expected 1:1 segregation.

- 5 The resistance had also introgressed into plants B23 and B28, as was shown in later generations.

Field test

- A preliminary field test was carried out with the resistant plants from the seedling test of the BC3 of B27. The plants were planted in a heavily infested field at our trial station at Kleve,
10 Germany. All plants stayed healthy and showed no symptoms when dug up at maturity.

Transfer to other *B. oleraceas*

The resistant B27 was used as source of the clubroot resistance for a backcross program in broccoli, cauliflower, cabbage and Brussels sprouts.

Example 2: Disease test

- 15 Seedlings are transplanted one week after sowing in multipots containing a mixture of sand and peatsoil at a rate of 2:1 with PH<6. One day after transplanting 1 ml of a solution of 1.10^6 cysts/ml is injected in the pots. The solution is of a mixture of clubroot sources with ECD codes 16/3/30, 16/23/30, 16/3/14 and 16/7/30 (Buczacki et al., 1975: Trans. Br. Mycol. Soc. 65, 295-303). The first two weeks the soil is kept wet, after that the soil may dry a bit.
20 Incubation temp is 18-20°C in the greenhouse. 4-5 weeks after transplanting the roots are washed clean and scored for the disease.

- Alternatively, seeds are directly sown in multipot trays in a standard sand : peat soil mixture (2:1 EGO, pH 5-6, 1 seed per multi-pot). Normally 20-30 plants/line are tested. Approx. 7-10 days after sowing the plants are inoculated, a second inoculation is followed 2-5 days after
25 the first inoculation. If seeds are suspected to germinate badly, sowing is done in black trays on rows and the plants are transplanted after 7-10 days to the multipot trays. Incubation temperature is 20-22 °C.

- Standard inoculum is isolate 9 (an aggressive isolate isolated from an infected field in Germany). The inoculum is prepared from infected roots stored in the fridge at -20 °C in
30 plastic bags. The roots are transferred to a mixer (1 part roots, 5 parts water) for ca. 2 min.

The small particles are then filtered with cheese cloth and the number of cysts (round, light (little blue) coloured) is counted. One ml of $0.5-1 \times 10^7$ cysts/ml is added to the stem/root at the base of the plant with a Eppendorf pipet. The inoculum needs to be shaken regularly to mix the available cysts in the suspension homogeneously.

- 5 Dependent on the size of the trial, sufficient controls need to be present. Susceptible controls are for example: White Rock (cauliflower), Maximus or other F1 (Brussels sprouts), Marathon (white cabbage). As resistant control Parkin or Storkin (Chinese cabbage) may be added. Hopkin (susceptible Chinese cabbage) may be added, as it gives in an early stage an indication if the inoculation has succeeded.
- 10 The plants need to be kept sufficiently moist during the trial. Regular fertilisation necessary, however plants must not grow excessively. Temperature is 22-20 °C (day/night), 16 hr light. After 4 (summer) – 6 (winter) weeks after inoculation plants can be pulled out of the soil, washed and scored for symptoms.

An observation scale for a disease test is for example:

- 15 0 = no clubs, healthy root system
- 1 = 1 - 2 small clubs on lateral roots, sometimes roots show some browning
- 2 = several small clubs on lateral roots, main root sometimes a little thickened
- 3 = main root thickened for largest part
- 4 = main root severely thickened and coalesced with clubbed lateral roots, some
- 20 normal roots still present
- 5 = one clump of roots, no normal roots present

- 25 The plants with score 0 (sometimes also the 1's, depending on the severity of the greenhouse test and the genetics involved) out of the greenhouse test are then transferred to a field infected with clubroot (e.g. field De Wit in Enkhuizen). Normally, escapes from the greenhouse become severely infected in the field and are readily visible (stunted growth). However, these are only very rare plants.

Another observation scale for a disease test is for example:

- 1 = No visible root galling
- 2 = Single gall on lateral roots
- 3 = Several small galls on lateral roots (plant healthy)

- 4 = Mild galling of the taproot, several small galls on lateral roots
- 5 = Moderate galling of the taproot, many small or several large galls on lateral roots
- 6 = Severe galling of the taproot, many large galls on laterals
- 7 = Severely galled, several healthy roots remaining
- 8 = Severly galled, few healthy roots present
- 9 = Severly galled, no healthy roots present

Example 3: Results of field trials

Table 1

Field trial in Werribee, Victoria, Australia. Plants transplanted in January 2002 to clubroot infested soil. An assessment after 6 weeks after transplantation shows (Table 1) already the effect of clubroot resistant cauliflower hybrids (F308 and F311) compared to susceptible varieties (White Rock and Triumphant) and a susceptible variety (Triumphant) treated with different amounts of a fungicide (Shirlane) and addition of lime (increasing pH heaving a reducing effect on clubroot severity, e.g. Dixon & Page, 1998, Acta Horticulturae, 459, 343-349). The observation scale above with rating from 1-9 was used in the trial below.

	Treatment	Cauli root score 0-9 scale
1	Quicklime (2.5t/ha)	4.11
2	Shirlan (3L/ha)	2.33
3	Shirlan (3L/ha) + Quicklime (2.5 t/ha)	2.22
4	Shirlan (2L/ha)	3.00
5	Control – Triumphant	5.06
6	F308	1.00
7	F311	1.00
8	White Rock	5.11

10

Table 2

Results from a young plant test of cauliflower with isolates originating from 3 different sites in Australia (Cora Lyn, Werribee and Trentham). The isolates were inoculated with a method comparable as described above. White Rock is a susceptible control and E70 is a resistant

hybrid cauliflower. The observation scale above with rating from 1-9 was used in the trial below.

Treatment averages			
	Cora Lyn	Werribee	Trentham
White Rock	8.2	8.7	3.9
E 70	1	1	1

Table 3

- 5 Results of field tests carried out at different locations in Europe naturally infested with clubroot in two years. D249, D506, E245, E246 are resistant cauliflower hybrids, White Rock is a susceptible cauliflower variety, SPR666 and A876 are resistant *B. sprouts* hybrid, Romulus and Maximus are susceptible *B. sprouts* varieties, F1182, F1187 are resistant white cabbage hybrids and Marathon a susceptible white cabbage variety. Assessments (di05) were
- 10 done at the end of the growing season on a 0-5 scale (see above).

entry	type	site	year	di05
D249	cauliflower	Hannover	2000	0.0
D506	cauliflower	Hannover	2000	0.0
White Rock	cauliflower	Hannover	2000	1.8
D249	cauliflower	Halsall-UK	2000	0.0
White Rock	cauliflower	Halsall-UK	2000	4.3
SPR666	B. sprouts	Halsall-UK	2000	0.0
Romulus	B. sprouts	Halsall-UK	2000	4.5
SPR666	B. sprouts	Roeselaere-Belgium	2000	0.0
Romulus	B. sprouts	Roeselaere-Belgium	2000	3.3
A876	B. sprouts	Halsall-UK	2001	0.0
Maximus	B. sprouts	Halsall-UK	2001	4.5
E245	cauliflower	Halsall-UK	2001	0.0
White Rock	cauliflower	Halsall-UK	2001	4.6
F1187	white cabbage	Halsall-UK	2001	0.0
Marathon	white cabbage	Halsall-UK	2001	3.5

E246	cauliflower	Hannover	2001	0.0
White Rock	cauliflower	Hannover	2001	1.9
F1182	white cabbage	Hannover	2001	0.0
Marathon	white cabbage	Hannover	2001	3.6

Example 4: Molecular markers

RAPDs markers linked to the resistance were developed using two white cabbage populations (D1544 and D1545). Two molecular markers obtained with primers O20 and Y13, respectively, resulted in PCR amplification products that are tightly linked to the resistance to clubroot. Both O20 and Y13 are 94.6% (53 out of 56) correlated with the result (resistant or sensitive) of disease test in population D1544, indicating a linkage distance of 5.4 cM. For population D1555, O20 is 100% while Y13 is 95.7% (45 out of 47) correlated with disease test, suggesting a tight linkage (0 cM) between O20 and the resistance and 4.3 cM between Y13 and the resistance. When testing the RAPDs in a dihaploid white cabbage population of 140 plants, O20 shows 100% (0 cM) and Y13 98.6% (1.4 cM) correlated with the result of disease tests.

Primer O20 (5'-ACA CAC GCT G-3') yields a specific fragment of approximately 400 bp.

Primer Y13 (5'-GGG TCT CGG T-3') yields a specific fragment of approximately 640 bp.

The amplification conditions are described below. After PCR 25 µl DNA is run on a 1.8% agarose gel.

- DNA	:	µl (diluted DNA from a standard miniprep)
- Primer (10 µM)	:	1.0 µl
- dNTP (2.5 mM)	:	2.0 µl
- Platinum Taq buffer 10x	:	2.5 µl (200mM Tris-HCl pH 8.4, 500 mM KCl minus MgCl ₂)
- MgCl ₂ (50 mM)	:	0.75 µl
- Platinum Taq (BRL/life)	:	0.2 µl (5 U/µl)
- sterile water:	:	µl (depends how much DNA is used)
25 µl final volume		

PCR program:

		3' 94°C		
	-	Ramp 0:00	94°C	0.10
5	-	Ramp 0:00	36°C	0.30
	-	Ramp 0:45	72°C	1.05
		3' 94°C		
	-	Ramp 0:00	94°C	0.10
		Ramp 0:00	36°C	0.30
10	-	Ramp 0:45	72°C	1.05
	-	5' 72°C		
				1 cycle, PCR cycle (Perkin Elmer 9600)
				40 cycles, PCR cycle (Perkin Elmer 9600)

SEQUENCE LISTING

<110> Syngenta Participations AG

5 <120> Disease resistant Brassicas

<130> 70059P

<140>

10 <141>

<160> 2

<170> PatentIn version 3.1

15

<210> 1

<211> 10

<212> DNA

<213> Artificial sequence

20

<220>

<223> Oligonucleotide

<400> 1

acacacgctg

10

25

<210> 2

<211> 10

<212> DNA

30 <213> Artificial sequence

<220>

<223> Oligonucleotide

<400> 2

35 gggctctcggt

10

CLAIMS

What is claimed is:

1. A *B. oleracea* plant resistant to clubroot, wherein the resistance to clubroot is
5 monogenic and dominant.
2. The plant according to claim 1, wherein said *B. oleracea* plant is rated at level 2 or less in a test for the disease having a 1-9 scale or at a level 1 or less in a test for the disease having a 0-5 scale.
3. The plant according to claim 1, wherein said *B. oleracea* plant is rated at level 1 in a
10 test for the disease having a 1-9 scale or at a level 0 in a test for the disease having a 0-5 scale.
4. The plant according to claim 1, wherein said *B. oleracea* plant is broccoli, white cabbage, cauliflower, Brussels sprouts, Borecole, Savoy, or red cabbage.
5. The plant according to claim 1, wherein said resistance is linked to a molecular
15 marker obtainable by PCR amplification using primer O20 (SEQ ID NO:1) or primer Y13 (SEQ ID NO:2).
6. The plant according to claim 5, wherein said resistance is within 10 cM of said molecular marker.
7. The plant according to claim 5, wherein said resistance is within 6 cM of said
20 molecular marker.
8. The plant according to claim 1, wherein said resistance is obtainable from a clubroot resistant *B. rapa* plant.
9. The plant according to claim 1, wherein said resistance is obtainable from Chinese cabbage F1 hybrid Parkin.
- 25 10. The plant according to any one of claims 1 to 9, wherein said plant is homozygous for said resistance.
11. The plant according to any one of claims 1 to 9, wherein said plant is heterozygous for said resistance.
12. The plant according to any one of claims 1 to 9, wherein said plant is an inbred.
- 30 13. The plant according to any one of claims 1 to 9, wherein said plant is a hybrid.

14. The plant according to claim 12 or 13, wherein said plant is cytoplasmic male sterile.
15. Seed of a plant according to any one of claims 1 to 14.
16. Use of a monogenic and dominant resistance to clubroot to confer a *B. oleracea* plant resistance to said disease, wherein said resistance is obtainable from a *B. rapa* plant.
- 5 17. Use of claim 16, wherein said resistance is obtainable from Chinese cabbage F1 hybrid Parkin.
18. A method for producing a *B. oleracea* plant comprising a monogenic and dominant resistance to clubroot comprising the steps of:
 - a) obtaining a *B. rapa* plant resistant to clubroot;
 - 10 b) crossing said *B. rapa* plant with a *B. oleracea* plant,
 - c) rescuing embryos resulting from the cross of step b);
 - d) regenerating a plant from an embryo of step c);
 - e) selecting a plant of step d) that is resistant to clubroot;
 - f) back-crossing a plant resulting from step e) with a *B. oleracea* plant.
- 15 19. The method according to claim 18, further comprising introgressing the resistance into an elite *B. oleracea* inbred.
20. The method according to claim 19, further comprising crossing said inbred to another *B. oleracea* inbred to produce a hybrid.
21. A *B. oleracea* plant obtainable by the method of any one of claims 18 to 20.
- 20 22. A method for transferring a monogenic and dominant resistance to clubroot to a *B. oleracea* plant susceptible or less resistant to the disease comprising the steps of:
 - a) obtaining a *B. oleracea* plant comprising a monogenic and dominant resistance to clubroot;
 - b) crossing said *B. oleracea* plant of step a) with a *B. oleracea* plant susceptible or less resistant to clubroot;
 - 25 c) selecting a plant from the cross of step b) that is resistant to clubroot.
23. The method according to claim 22, further comprising backcrossing said resistance into said *B. oleracea* plant susceptible or less resistant to clubroot.
24. A DNA fragment amplified from a Brassica genome, wherein said DNA fragment is approximately 400 bp long and comprises SEQ ID NO:1.
- 30 25. A DNA fragment amplified from a Brassica genome, wherein said DNA fragment is approximately 640 bp long and comprises SEQ ID NO:2.

26. A DNA fragment according to claim 24 or 25, wherein said DNA fragment is indicative of the presence of a dominant and monogenic resistance to clubroot in a Brassica plant.
27. Use of a DNA fragment according to claim 24 or 25 to identify a Brassica plant that is resistant to clubroot.
28. Use of primer O20 (SEQ ID NO:1) to detect a DNA fragment of approximately 400 bp in a Brassica genome.
29. Use of primer Y13 (SEQ ID NO:2) to detect a DNA fragment of approximately 640 bp in a Brassica genome.
30. Use of a primer O20 or Y13 to identify a Brassica plant resistant to clubroot.
31. Use of any one of claims 27-30, wherein said Brassica plant is *B. oleracea*.
32. A kit for detecting a monogenic and dominant resistance to clubroot in a *B. oleracea* plant comprising an oligonucleotide of set forth in SEQ ID NO:1 or SEQ ID NO:2.
33. A method for transferring a monogenic and dominant resistance to clubroot to a *B. oleracea* plant susceptible or less resistant to the disease comprising the steps of:
- obtaining a *B. oleracea* plant comprising a monogenic and dominant resistance to clubroot;
 - crossing said *B. oleracea* plant of step a) with a *B. oleracea* plant susceptible or less resistant to clubroot;
 - selecting a plant comprising a DNA fragment obtainable by PCR amplification using primer O20 (SEQ ID NO:1) or primer Y13 (SEQ ID NO:2);
- wherein said plant of step c) is resistant to clubroot.
34. The method according to claim 33, further comprising backcrossing said resistance into said *B. oleracea* plant susceptible or less resistant to clubroot.
35. A method of identifying a *B. oleracea* plant resistant to clubroot comprising the steps of:
- obtaining a sample from a *B. oleracea* plant;
 - detecting in said sample a DNA fragment obtainable by PCR amplification using primer O20 (SEQ ID NO:1) or primer Y13 (SEQ ID NO:2),
- wherein said *B. oleracea* plant of step b) is resistant to clubroot.

36. A method of selecting a *B. oleracea* plant resistant to clubroot from a population of *B. oleracea* plants comprising the steps of:

- a) providing a population of *B. oleracea* plants;
- b) obtaining a sample of a plant of said population;
- c) detecting in said sample a DNA fragment obtainable by PCR amplification

5

using primer O20 (SEQ ID NO:1) or primer Y13 (SEQ ID NO:2),
wherein said *B. oleracea* plant of step b) is resistant to clubroot.

ABSTRACT

DISEASE RESISTANT BRASSICAS

5 The present invention discloses *B. oleracea* plants resistant to clubroot disease. In particular, the plants of the present invention comprise a monogenic dominant resistance to the disease clubroot introgressed from *B. rapa*. This resistance provides improved resistance to the disease as compared to previously existing resistances in *B. oleracea*.